ELSEVIER

Journal of Organometallic Chemistry 539 (1997) 209

Journal of Organometallic Chemistry

Author Index of Volume 539

Akita, M., 205
Alekseev, V.I., 27
Aoki, Y., 115
Aroney, M.J., 121

Baguley, P., 193
Barnes, C.M., 121
Bényei, A.C., 177
Bosque, R., 99
Bruce, M.I., 141

Camalli, M., 9
Campi, E.M., 147
Castellanos-Páez, A., 1
Castillón, S., 1
Cecchi, P., 9
Claver, C., 1
Cooks, R.G., 131

Dias, H.V.R., 77
Digilio, G., 9
Donaldson, W.A., 87
Duesler, E.N., 19, 37

Ferrand, V., 201
Font-Bardía, M., 99
Fukui, K., 115

Gabbai, F.P., 187
Gau, H.-M., 155
Gioia Lobbia, G., 9
Glidewell, C., 177
Gobetto, R., 9
Guénin, E., 193

Heidel, H., 67
Hoffmann, A., 61
Hor, T.S.A., 45
Hua, R., 205
Hughes, A.K., 109
Humphrey, P.A., 141
Huttner, G., 67

Ikeda, S., 115

Jackson, W.R., 147
Janik, J.F., 19
Jia, G., 53
Jiang, M.-K., 155
Jin, P., 171

Kamigata, N., 171
Kaul, F.A.R., 187
Kingsley, A.J., 109
Kirin, V.P., 27
Kitagawa, S., 171
Kondo, M., 171

Le Bozec, H., 201
Leininger, S., 61
Lightfoot, P., 177
Lin, C.-C., 155
Liu, Z., 127
López, C., 99
Low, P.M.N., 45

Ma, S., 131
Mak, T.C.W., 45, 127
Maksakov, V.A., 27
McCubbin, Q.J., 147
Moro-oka, Y., 205
Muir, K.W., 193

Paine, R.T., 19, 37
Péron, V., 201
Pétillon, F.Y., 193
Phillips, L., 121
Plasseraud, L., 163
Podberezskaya, N.V., 27
Poder-Guillou, S., 193
Porhiel, E., 201

Ramaswamy, M., 87
Regitz, M., 61
Reisacher, H.-U., 37
Royles, B.J.L., 177

Scherer, J., 67
Schollhammer, P., 193
Shang, L., 87
Shimizu, T., 171
Skelton, B.W., 141
Smith, D.M., 177
Solans, X., 99
Spagna, R., 9
Süss-Fink, G., 163

Talarmin, J., 193
Tao, C., 87
Tkachev, S.V., 27
Trnacek, A.E., 147
Tschinkl, M., 187

Urakubo, T., 171

Virovets, A.V., 27

Wang, F., 131
Wang, Z., 77
White, A.H., 141
Wong, P.S.H., 131
Wu, W.F., 53

Xia, H.P., 53
Xie, Z., 127
Xue, F., 127

Yamada, I., 115
Yamagishi, T., 115
Yamaguchi, M., 115
Yeung, R.C.Y., 53
Young Jr., V.G., 87
Yun, Y.K., 87

Zhang, Z.-Y., 45

Elsevier Science S.A.

ELSEVIER

Journal of Organometallic Chemistry 539 (1997) 211–215

Journal of Organometallic Chemistry

Subject Index of Volume 539

Acyl ferrocenes
Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1′-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177
Allylic cleavage
Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147
Aluminium
Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkl and F.P. Gabbai), 187
Preparation and characterization of bis-(trimethylsilyl)aminoalanes $[H_2AlN(SiMe_3)_2] \cdot NMe_3$ and $\{HAl[N(SiMe_3)_2]_2\}_n$. Crystal structure determination for tris-(trimethylsilyl)alane trimethylamine adduct $(Me_3Si)_3Al \cdot NMe_3$ (J.F. Janik, E.N. Duesler and R.T. Paine), 19
Amines
Preparation and characterization of bis-(trimethylsilyl)aminoalanes $[H_2AlN(SiMe_3)_2] \cdot NMe_3$ and $\{HAl[N(SiMe_3)_2]_2\}_n$. Crystal structure determination for tris-(trimethylsilyl)alane trimethylamine adduct $(Me_3Si)_3Al \cdot NMe_3$ (J.F. Janik, E.N. Duesler and R.T. Paine), 19
Aminoalkenylcarbene
Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201
4-Aminobenzoylferrocene
Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1′-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177
Anion–cation interaction
Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171
Aromatics
Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163
Asymmetric hydroformylation
Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptomethyl)-bicyclo[2.2.2]octane (H_2-BCOS). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1
Asymmetric hydrogenation
Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115
1,4-Asymmetric induction
Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115
Benzenechromiumtricarbonyl
The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121
Bifunctional Lewis acids
Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkl and F.P. Gabbai), 187
Bimetallic complexes
The synthesis and characterization of Group 4 metal–chromium complexes bridged by an $OCH_2C_5H_4$ group. The molecular structure of $Cp_2TiCl[(OCH_2C_5H_4)Cr(CO)_2(NO)]$ (M.-K. Jiang, C.-C. Lin and H.-M. Gau), 155
Biphasic media
Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163
Bis(carbene)
Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201
Bond moment
The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121
Boron
Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano–mercury(II) polypyrazolylborates. X-ray crystal structure of $[HB(3,5\text{-}Me_2pz)_3]HgCN$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9
Carbenoid
Reactions of a tungsten phosphenium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37
Carbonyl clusters
Syntheses and characterization of $OsPt_2(\mu\text{-}CO)(\mu\text{-}I)_2(CO)_2(PPh_3)_3$ and $Os_2(\mu\text{-}I)_2(CO)_{6-n}(PPh_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podberezskaya), 27
Catalysis
Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147

Elsevier Science S.A.

Chiral phosphine ligands
Rhodium–COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67
Chiral phosphite ligands
Rhodium–COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67
Chiral rhodium complexes
Rhodium–COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67
Chiral thiolate
Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptomethyl)-bicyclo[2.2.2]octane (H_2-BCOS). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1
Chiral tripodal ligands
Rhodium–COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67
Chromium charge
The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121
Cluster
Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163
Cobalt
Stereoelectronic effects and gas phase Co^+, Ni^+, $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131
π-Conjugated bridges
Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201
Cooligomerization
Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61
Crystal structure
Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171
Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1′-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-}Cl)(\mu\text{-}SMe)_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-}SCH_2CH_2SH)(\mu\text{-}SMe)_3]\cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Cyclisation
Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147
Cyclopentadienyl
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-}Cl)(\mu\text{-}SMe)_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-}SCH_2CH_2SH)(\mu\text{-}SMe)_3]\cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Synthesis and crystal structure of an organolanthanide fluoride, $[\{(Me_3Si)_2C_5H_3\}_2Sm(\mu\text{-}F)]_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127
Cyclopentadienyl groups
Syntheses and characterization of zirconium and hafnium complexes of amido-fluorenyl ligands $[(NBu^t)SiMe_2CH_2(C_{13}H_8)]^{2-}$ and $[(NPr^i)SiMe_2CH_2(C_{13}H_8)]^{2-}$ (H.V.R. Dias and Z. Wang), 77
Cyclopentadienyls
Reactions of a tungsten phosphenium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37
Stereoelectronic effects and gas phase Co^+, Ni^+, $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131
Dehydrodipeptide
Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115
Dephenylation
Trapping of a phenyl group by a ruthenium cluster during dephenylation of coordinated $CH_2(PPh_2)_2$ (dppm) (M.I. Bruce, P.A. Humphrey, B.W. Skelton and A.H. White), 141
Diene complexes
Reactivity of 1-substituted (pentadienyl)iron(1+) cations: regioselectivity for addition of malonate nucleophiles; formation of (pentenediyl)- and (diene)iron complexes (W.A. Donaldson, L. Shang, C. Tao, Y.K. Yun, M. Ramaswamy and V.G. Young Jr.), 87
Dimers
Dimeric and polymeric ruthenium complexes with Ru–vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53
Dinuclear complex
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-}Cl)(\mu\text{-}SMe)_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-}SCH_2CH_2SH)(\mu\text{-}SMe)_3]\cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Dithiol
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-}Cl)(\mu\text{-}SMe)_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-}SCH_2CH_2SH)(\mu\text{-}SMe)_3]\cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Electrostatic interaction
Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115
Enantioselective catalysis
Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptomethyl)-bicyclo[2.2.2]octane (H_2-BCOS). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1
Enantioselective hydrogenation
Rhodium–COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67

EXAFS
The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121
Ferrocenyloximes
Palladium(II) compounds containing σ[Pd–C_{sp^2}(ferrocene)] bonds and ferrocenyloximes as bidentate {C N}– ligands. X-ray crystal structure of [Pd{[(η^5-C_5H_3)–C(CH_3)=N(OH)]Fe(η^5-C_5H_5)}-Cl(PPh_3)] (C. López, R. Bosque, X. Solans and M. Font-Bardía), 99
Fluorine
Synthesis and crystal structure of an organolanthanide fluoride, [{(Me_3Si)$_2$$C_5H_3$}$_2$Sm($\mu$-F)]$_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127
Halide
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex [Cp_2Mo_2(μ-Cl)(μ-SMe)$_3$]: crystal structure of [Cp_2Mo_2(μ-SCH_2CH_2SH)(μ-SMe)$_3$] · 0.5CH_2Cl_2 (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Hydrides
Preparation and characterization of bis-(trimethylsilyl)aminoalanes [H_2AlN($SiMe_3$)$_2$] · NMe_3 and {HAl[N($SiMe_3$)$_2$]$_2$}$_n$. Crystal structure determination for tris-(trimethylsilyl)alane trimethylamine adduct (Me_3Si)$_3$Al · NMe_3 (J.F. Janik, E.N. Duesler and R.T. Paine), 19
Hydrochinone
Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkl and F.P. Gabbai), 187
Hydrogenation
Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163
4-Hydroxybenzoylferrocene
Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1′-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177
1,1′-bis(4-Hydroxybenzoyl)ferrocene
Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1′-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177
Induced fitting
Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115
Infrared force constants
The charge distribution in substituted benzenechromiumtricarbonyls: an interpretation of spectroscopic data in terms of bond dipoles and charges (L. Phillips, C.M. Barnes and M.J. Aroney), 121
Iron
Reactivity of 1-substituted (pentadienyl)iron(1 +) cations: regioselectivity for addition of malonate nucleophiles; formation of (pentenediyl)- and (diene)iron complexes (W.A. Donaldson, L. Shang, C. Tao, Y.K. Yun, M. Ramaswamy and V.G. Young Jr.), 87
Stereoelectronic effects and gas phase Co^+, Ni^+, $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131
Lanthanide
Synthesis and crystal structure of an organolanthanide fluoride, [{(Me_3Si)$_2$$C_5H_3$}$_2$Sm($\mu$-F)]$_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127
Ligand exchange
Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163
Mass spectrometry
Stereoelectronic effects and gas phase Co^+, Ni^+, $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131
Mercury
Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano–mercury(II) polypyrazolylborates. X-ray crystal structure of [HB(3,5-Me_2pz)$_3$]HgCN (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9
Metal–carbon bonds
Titanium and zirconium neopentyl chloro complexes, MNp_xCl_{4-x} (x = 1–4) (A.K. Hughes and A.J. Kingsley), 109
Metal–chromium complexes
The synthesis and characterization of Group 4 metal–chromium complexes bridged by an $OCH_2C_5H_4$ group. The molecular structure of Cp_2TiCl[($OCH_2C_5H_4$)Cr($CO)_2$(NO)] (M.-K. Jiang, C.-C. Lin and H.-M. Gau), 155
Metallo-phosphane complex
Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61
Metathesis
General synthesis of ferrocenylphosphine-bridged digold carboxylato complexes. Crystal and molecular structure of Au_2(η^1-CF_3CO_2)$_2$-(μ-dppf) · C_6H_{14} (dppf = Fe($C_5H_4PPh_2$)$_2$) (P.M.N. Low, Z.-Y. Zhang, T.C.W. Mak and T.S.A. Hor), 45
Methoxyalkenylcarbene
Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201
Molecular structure
Functionalised acyl ferrocenes: crystal and molecular structures of 4-aminobenzoylferrocene, 4-hydroxybenzoylferrocene and 1,1′-bis(4-hydroxybenzoyl)ferrocene (A.C. Bényei, C. Glidewell, P. Lightfoot, B.J.L. Royles and D.M. Smith), 177
Palladium(II) compounds containing σ[Pd–C_{sp^2}(ferrocene)] bonds and ferrocenyloximes as bidentate {C N}– ligands. X-ray crystal structure of [Pd{[(η^5-C_5H_3)–C(CH_3)=N(OH)]Fe(η^5-C_5H_5)}-Cl(PPh_3)] (C. López, R. Bosque, X. Solans and M. Font-Bardía), 99
Trapping of a phenyl group by a ruthenium cluster during dephenylation of coordinated $CH_2(PPh_2)_2$ (dppm) (M.I. Bruce, P.A. Humphrey, B.W. Skelton and A.H. White), 141
Molybdenum
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex [Cp_2Mo_2(μ-Cl)(μ-SMe)$_3$]: crystal structure of [Cp_2Mo_2(μ-SCH_2CH_2SH)(μ-SMe)$_3$] · 0.5CH_2Cl_2 (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Monophosphines
General synthesis of ferrocenylphosphine-bridged digold carboxylato complexes. Crystal and molecular structure of Au_2(η^1-CF_3CO_2)$_2$-(μ-dppf) · C_6H_{14} (dppf = Fe($C_5H_4PPh_2$)$_2$) (P.M.N. Low, Z.-Y. Zhang, T.C.W. Mak and T.S.A. Hor), 45

NMR spectroscopy
Syntheses and characterization of zirconium and hafnium complexes of amido-fluorenyl ligands $[(NBu^t)SiMe_2CH_2(C_{13}H_8)]^{2-}$ and $[(NPr^i)SiMe_2CH_2(C_{13}H_8)]^{2-}$ (H.V.R. Dias and Z. Wang), 77
Optically active compound
Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171
Osmium
Syntheses and characterization of $OsPt_2(\mu\text{-}CO)(\mu\text{-}I)_2(CO)_2(PPh_3)_3$ and $Os_2(\mu\text{-}I)_2(CO)_{6-n}(PPh_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podberezskaya), 27
Palladium(II) compounds
Palladium(II) compounds containing σ[Pd–C_{sp^2}(ferrocene)] bonds and ferrocenyloximes as bidentate {C N}– ligands. X-ray crystal structure of $[Pd\{[(\eta^5\text{-}C_5H_3)\text{–}C(CH_3)\text{=}N(OH)]Fe(\eta^5\text{-}C_5H_5)\}Cl(PPh_3)]$ (C. López, R. Bosque, X. Solans and M. Font-Bardía), 99
Phosphaalkyne cyclooligomers
Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61
Phosphaalkynes
Organophosphorus compounds 115.. Solvent-directed cooligomerization of kinetically stabilized phosphaalkynes with trialkylaluminium compounds (A. Hoffmann, S. Leininger and M. Regitz), 61
Phosphenium ion
Reactions of a tungsten phosphenium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37
Phosphine
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-}Cl)(\mu\text{-}SMe)_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-}SCH_2CH_2SH)(\mu\text{-}SMe)_3]\cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Phosphine bridges
General synthesis of ferrocenylphosphine-bridged digold carboxylato complexes. Crystal and molecular structure of $Au_2(\eta^1\text{-}CF_3CO_2)_2(\mu\text{-dppf})\cdot C_6H_{14}$ (dppf = $Fe(C_5H_4PPh_2)_2$) (P.M.N. Low, Z.-Y. Zhang, T.C.W. Mak and T.S.A. Hor), 45
Phosphorus
Reactions of a tungsten phosphenium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37
Platinum
Syntheses and characterization of $OsPt_2(\mu\text{-}CO)(\mu\text{-}I)_2(CO)_2(PPh_3)_3$ and $Os_2(\mu\text{-}I)_2(CO)_{6-n}(PPh_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podberezskaya), 27
Polymers
Dimeric and polymeric ruthenium complexes with Ru–vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53
Poly(pyrazolyl)borates
Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano–mercury(II) polypyrazolylborates. X-ray crystal structure of $[HB(3,5\text{-}Me_2pz)_3]HgCN$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9
Proton affinities
Stereoelectronic effects and gas phase Co^+, Ni^+, $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131
Pyridine
Stereoelectronic effects and gas phase Co^+, Ni^+, $CpFe^+$, $CpCo^+$ and $CpNi^+$ affinities of pyridines studied by the kinetic method (P.S.H. Wong, S. Ma, F. Wang and R.G. Cooks), 131
Rearrangement
Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147
Resorcinol
Alumination of resorcinol and hydroquinone. An easy access to bifunctional Lewis acids (F.A.R. Kaul, M. Tschinkl and F.P. Gabbai), 187
Rhodium
Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147
Rhodium catalyst
Electrostatic interaction and induced fitting of the rhodium(I) complex coordinated by diphosphine ligand having an amino group in the diastereoselective hydrogenation of dehydrodipeptides (I. Yamada, K. Fukui, Y. Aoki, S. Ikeda, M. Yamaguchi and T. Yamagishi), 115
Rhodium COD complexes
Rhodium–COD-Komplexe chiraler Tripod-Liganden mit drei verschiedenen Donorgruppen: Komplexchemie und Katalyse (J. Scherer, G. Huttner und H. Heidel), 67
Rhodium complexes
Synthesis of rhodium(I) complexes with the new dithiol chiral ligand (+)-*trans*-2,3-bis(mercaptomethyl)-bicyclo[2.2.2]octane (H_2-BCOS). Their application as catalysts precursors in the styrene hydroformylation (A. Castellanos-Páez, S. Castillón and C. Claver), 1
Ruthenium
Catalytic hydrogenation of benzene derivatives under biphasic conditions (L. Plasseraud and G. Süss-Fink), 163
Dimeric and polymeric ruthenium complexes with Ru–vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53
Ruthenium cluster
Trapping of a phenyl group by a ruthenium cluster during dephenylation of coordinated $CH_2(PPh_2)_2$ (dppm) (M.I. Bruce, P.A. Humphrey, B.W. Skelton and A.H. White), 141
Samarium
Synthesis and crystal structure of an organolanthanide fluoride, $[\{(Me_3Si)_2C_5H_3\}_2Sm(\mu\text{-}F)]_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127
Selenonium salt
Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171
Silicon
Preparation and characterization of bis-(trimethylsilyl)aminoalanes $[H_2AlN(SiMe_3)_2]\cdot NMe_3$ and $\{HAl[N(SiMe_3)_2]_2\}_n$. Crystal structure determination for tris-(trimethylsilyl)alane trimethylamine adduct $(Me_3Si)_3Al\cdot NMe_3$ (J.F. Janik, E.N. Duesler and R.T. Paine), 19
Solid state ^{15}N NMR
Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano–mercury(II) polypyrazolylborates. X-ray crystal structure of $[HB(3,5\text{-}Me_2pz)_3]HgCN$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9

Solid-state structure
Syntheses and characterization of $OsPt_2(\mu\text{-CO})(\mu\text{-I})_2(CO)_2(PPh_3)_3$ and $Os_2(\mu\text{-I})_2(CO)_{6-n}(PPh_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podberezskaya), 27
Solution structure
Syntheses and characterization of $OsPt_2(\mu\text{-CO})(\mu\text{-I})_2(CO)_2(PPh_3)_3$ and $Os_2(\mu\text{-I})_2(CO)_{6-n}(PPh_3)_n$ ($n = 1, 2$) in the solid-state and solution (V.A. Maksakov, V.P. Kirin, A.V. Virovets, S.V. Tkachev, V.I. Alekseev and N.V. Podberezskaya), 27
Stoichiometric complexes
Preparation and reactions of rhodium complexes of some alkenyloxy- and alkenylamino-benzylamines and related compounds: a comparison with catalytic reactions (E.M. Campi, W.R. Jackson, Q.J. McCubbin and A.E. Trnacek), 147
Telluronium salt
Crystal structures of optically active diastereomeric telluronium and selenonium salts. Anion–cation interactions in the crystalline state (T. Shimizu, T. Urakubo, P. Jin, M. Kondo, S. Kitagawa and N. Kamigata), 171
Thiolate
Reactions of dinuclear and polynuclear complexes. Part 18. Substitution reactions of a μ-chloro or a μ-thiolato ligand in the dinuclear cyclopentadienyl molybdenum(III) complex $[Cp_2Mo_2(\mu\text{-Cl})(\mu\text{-SMe})_3]$: crystal structure of $[Cp_2Mo_2(\mu\text{-SCH}_2CH_2SH)(\mu\text{-SMe})_3] \cdot 0.5CH_2Cl_2$ (P. Schollhammer, E. Guénin, S. Poder-Guillou, F.Y. Pétillon, J. Talarmin, K.W. Muir and P. Baguley), 193
Titanium
Titanium and zirconium neopentyl chloro complexes, MNp_xCl_{4-x} ($x = 1$–4) (A.K. Hughes and A.J. Kingsley), 109
Trimethylsilyl
Synthesis and crystal structure of an organolanthanide fluoride, $[\{(Me_3Si)_2C_5H_3\}_2Sm(\mu\text{-F})]_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127
Tungsten
Reactions of a tungsten phosphenium ion complex with carbenoid reagents (H.-U. Reisacher, E.N. Duesler and R.T. Paine), 37
Synthesis of mixed bis(alkenylcarbene) ditungsten complexes by sequential activation of dialkynols (V. Péron, E. Porhiel, V. Ferrand and H. Le Bozec), 201
Vinyl ruthenium compounds
Dimeric and polymeric ruthenium complexes with Ru–vinyl linkages (G. Jia, W.F. Wu, R.C.Y. Yeung and H.P. Xia), 53
X-ray
Metal polypyrazolylborates. XIII. Solution and solid state NMR study of cyano–mercury(II) polypyrazolylborates. X-ray crystal structure of $[HB(3,5\text{-}Me_2pz)_3]HgCN$ (G. Gioia Lobbia, P. Cecchi, R. Gobetto, G. Digilio, R. Spagna and M. Camalli), 9
X-ray diffraction
Synthesis and crystal structure of an organolanthanide fluoride, $[\{(Me_3Si)_2C_5H_3\}_2Sm(\mu\text{-F})]_2$ (Z. Xie, Z. Liu, F. Xue and T.C.W. Mak), 127
Zirconium
Syntheses and characterization of zirconium and hafnium complexes of amido-fluorenyl ligands $[(NBu^t)SiMe_2CH_2(C_{13}H_8)]^{2-}$ and $[(NPr^i)SiMe_2CH_2(C_{13}H_8)]^{2-}$ (H.V.R. Dias and Z. Wang), 77
Titanium and zirconium neopentyl chloro complexes, MNp_xCl_{4-x} ($x = 1$–4) (A.K. Hughes and A.J. Kingsley), 109